

Case Report

# Symmetry of Shank Muscle Strength, Passive Stiffness and Plantar Pressure Following IASTM Accompanied by Electrotherapy in a Case with Severe Ankle Stiffness

Fateme Khorramroo, Seyed Hamed Mousavi\* and Hooman Minoonejad

Faculty of Sport Science and Health, University of Tehran, Iran

## Summary

Excessive ankle stiffness can greatly impact mobility, leading to discomfort, difficulty in walking, and limited Range of Motion (ROM). We aimed to identify and address the symmetry of shank muscle strength, ankle passive stiffness, and plantar pressure distribution, in a patient with unilateral excessive ankle stiffness, utilizing Instrument-Assisted Soft Tissue Mobilization (IASTM) accompanied by Faradic Electrical Stimulation (FES).

The patient's muscle strength and ROM which had diminished due to 3.5 years of ankle immobilization post-rescue from amputation, underwent a 12-week program involving IASTM and FES. The plantar and dorsiflexion muscles' torque, ROM, and plantar pressure were measured using an isokinetic and plantar distribution system before and after the intervention. Symmetry of muscle torque, ROM, and plantar pressure between two limbs were calculated for pre and post-test.

Results indicated improvement in the ratio index of the concentric/eccentric dorsi- and plantar-flexion peak torque and dorsi- and plantar-flexion work, ROM, gait line length, and contact time after a 12-week intervention.

The study suggests that IASTM and FES are effective interventions for restoring symmetry in a patient with post-operation complications, highlighting the need for further research on similar cases.

## Introduction

Ankle fractures are relatively frequent injuries [1]. Although patients typically anticipate a full return to their pre-injury activities, many still encounter stiffness that can impede both their daily functions and overall recovery [2]. Ankle stiffness can significantly impair mobility and quality of life, leading to discomfort and functional limitations during daily activities [3]. Moreover, lower limb malalignment can lead to several issues including low back pain [4].

Severe ankle stiffness, often a consequence of prolonged immobilization or surgical interventions, can restrict the Range of Motion (ROM) [5] and alter muscle strength distribution in the affected limb [6]. This condition is particularly challenging in patients recovering from traumatic injuries or surgeries, where the balance of muscle strength between limbs becomes crucial for decreasing injury risk [7].

In cases of unilateral excessive ankle stiffness, discrepancies in muscle strength and passive stiffness between the affected and healthy limbs can exacerbate mobility issues, making it essential to explore effective rehabilitation strategies [5]. Not conventional treatments nor cyclic interventions including gait retraining [8,9] or specific shoes [10] may not fully address the underlying biomechanical imbalances and asymmetry, necessitating the consideration of specialized interventions.

This study focuses on a patient with a history of severe ankle stiffness following 3.5 years of immobilization due to post-op complications. The patient underwent a 12-week rehabilitation program incorporating Instrument-Assisted Soft Tissue Mobilization (IASTM) and Faradic Electrical Stimulation (FES). Our approach aimed to restore symmetry between two limbs in muscle strength, passive stiffness, and plantar pressure distribution during walking. By objectively measuring muscle torque, ROM, and plantar pressure before

## More Information

### \*Address for correspondence:

Seyed Hamed Mousavi, Faculty of Sport Science and Health, University of Tehran, Iran,  
Email: musavihamed@ut.ac.ir

 <https://orcid.org/0000-0003-2906-2101>

**Submitted:** November 09, 2024

**Approved:** November 13, 2024

**Published:** November 14, 2024

**How to cite this article:** Khorramroo F, Mousavi SH, Minoonejad H. Symmetry of Shank Muscle Strength, Passive Stiffness and Plantar Pressure Following IASTM Accompanied by Electrotherapy in a Case with Severe Ankle Stiffness. *J Sports Med Ther.* 2024; 9(4): 091-095. Available from: <https://dx.doi.org/10.29328/journal.jsmt.1001089>

**Copyright license:** © 2024 Khorramroo F, et al.

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Keywords:** Contact time; Path length; Torque; Ratio index



and after the intervention, we aimed to assess the effectiveness of IASTM and FES in addressing post-operative complications associated with severe ankle stiffness. The study's findings may provide valuable insights into rehabilitation practices for similar cases, highlighting the importance of restoring symmetry for enhanced mobility and functional recovery.

## Methods

### Case description

A 41-year-old female athlete has been engaging in regular exercise since the age of 10. At the age of 34, the patient experienced a severe injury to her right shank, which involved fractures of the tibia, fibula, calcaneus, and rupture of the Achilles tendon. Following the injury, the patient underwent several surgeries and experienced various complications.

The patient [11] was able to walk after 3.5 years (age: 37.5). It has been 3.5 years that the patient has had a normal life and frequent workouts (age: 41, at the time of the study). The patient had 2 cm of leg length discrepancy due to reduced heel fat pad and ankle joint interarticular space, reduced muscle strength, and ankle dorsiflexion ROM. Thus, the participant uses an orthosis under the right heel. Also, the participant had stiff ankles and Metatarsophalangeals (MTPs) in all dimensions. Participant continued to perform daily workouts including swimming and fitness training and did not make any change in exercise routine. The effect of interventions was assessed from Dec 2022 until Mar 2023 and this paper is a part of a research project [12].

The experiments reported in the manuscript were performed in accordance with the ethical standards of the Helsinki Declaration. This study was approved by the Research Ethics Committees of the Faculty of Physical Education and Sport Sciences-Tehran University (IR.UT.SPORT.REC.1401.045). The participant signed a written consent form.

### Plantar pressure path length and area

Single limb standing was measured while the participant stood on a foot pressure instrument (FDM-S; Zebris Medical GmbH, Weitnau-Seltmans, Germany) with the other limb in 30° of hip flexion and 45° of knee flexion and hands on hips. The participant was instructed to “stand as still as possible”. The participant performed 3 practice trials followed by three successful 10-second trials with eyes open and one minute rest between trials recorded. Unsuccessful trials, where the participant deviated from the test position, were repeated [13].

### Isokinetic measurements

The foot of the test limb was firmly strapped to the dynamometer foot-plate. The lateral malleolus was aligned with the axis of rotation of the dynamometer head and, using the lateral femoral condyle and the greater trochanter,

the knee was positioned in a statically flexed position (10°; goniometer).

### Passive dorsiflexion stiffness

The non-weight-bearing dorsiflexion ROM and passive dorsiflexion stiffness were calculated by measuring the slope of the torque-angle curve generated for 1° to 6° dorsiflexion [14].

### Isokinetic plantarflexion and dorsiflexion test

The isokinetic plantarflexion and dorsiflexion testing was performed at 180 °/s (the only possible velocity for the participant) with a sampling frequency of 1000 Hz. During test contractions, the participant was instructed to move her ankle through her full ROM as fast as possible and to push and pull as hard as possible in each direction. Strong verbal encouragement was provided throughout testing [15]. In order to maintain consistent contraction durations, the participant executed 10 isokinetic repetitions.

### Intervention

Interventions [16] were delivered by a registered physiotherapist (HM) and instructor (FKH). The program consisted of 3 sessions a week for 12 weeks; the participant warms up for 5 minutes, moving the ankle in 3 planes of movement. Two interventions were prescribed i.e., IASTM [17] techniques based on the manual for Graston technic [18] and FES [19]. The interventions were applied on separate days alternatively. Alternately, one-day electrotherapy on tibialis anterior, gastrocnemius muscles and foot soles, and Myofascial release [20] Graston technique (using GT5, GT6, and GT2) [21] friction massage [18] on calf muscles, Achilles tendon, sole of the foot and on the spots with adhesions on the other day. Electrotherapy [22] was used on the same day as participants' personal exercise. Beurer EM 49 Digital TENS/EMS was used for electrotherapy. The impulse was 250 μs duration and the frequency was 25 to 50 Hz. The time of each intervention was increased by 2-3 minutes every two weeks, whenever the participant was ready and reported no pain or inconvenience after each first session of increased time and pressure.

### Data analysis

Ratio index [17], uses the ratio of the two limbs values as the index of symmetry. For a variable X, it is defined as Ratio index =  $(1 - X_R/X_L)$ . 100%. Ratio index = 0% indicates full symmetry, while ratio index > 100% indicates asymmetry. This is a case study, and inferential statistics have not been performed for data analysis. Only descriptive statistics have been used

## Results

The participant is a 41-year-old individual with a height of 1.68 m and a body mass of 59 kg. The BMI is calculated



at 20.9, with leg lengths measuring 84.5 cm for the right leg and 86.5 cm for the left leg. Table 1 presents the results of the symmetry of peak torque, work, and ROM between two limbs. Table 2 shows the symmetry for the area of the center of pressure trajectory and path length. Table 3 illustrates results of symmetry for ankle passive stiffness in the sagittal plane and Table 4 shows the results of symmetry for gait line length and contact time. The value of ratio index = 0 indicates full symmetry.

The results showed an improvement in the ratio index of the concentric/eccentric dorsi- and plantar-flexion peak torque, ROM, and gait line length, Ratio Index of eccentric/concentric dorsi- and plantar-flexion work and contact time after a 12-week intervention.

## Discussion

The findings of this study highlight the significant impact of a 12-week rehabilitation program utilizing IASTM FES on restoring symmetry in muscle strength, passive stiffness, and plantar pressure distribution in a patient with unilateral excessive ankle stiffness. The results indicate that both interventions were effective in improving the ratio index of peak torque, ROM, and gait parameters, suggesting a positive shift towards normalizing biomechanical function.

### Symmetry in muscle strength and passive stiffness

The restoration of symmetry in muscle strength between the affected and unaffected limbs is critical for optimal mobility and functional recovery [23]. The improvement in

the ratio index for concentric and eccentric dorsi- and plantar-flexion peak torque demonstrates that the rehabilitation program successfully enhanced muscle performance on the previously affected side. This is particularly important given that prolonged immobilization often leads to muscle atrophy and strength deficits, which can exacerbate functional limitations [24]. The observed increase in symmetry of passive stiffness suggests a potential improvement in the mechanical properties of the ankle joint, which could facilitate better movement patterns during daily activities.

### Symmetry in plantar pressure distribution

Changes in plantar pressure distribution are indicative of altered gait mechanics and weight-bearing strategies. The study's results showed improvements in both the area of the center of pressure trajectory and path length, which are essential for maintaining balance and stability during ambulation. These findings suggest that the interventions not only addressed strength deficits but also contributed to more efficient weight transfer which is often affected in patients with unilateral ankle stiffness.

The successful application of IASTM and FES in this case underscores their potential as effective therapeutic modalities for addressing post-operative complications associated with severe ankle stiffness. The individualized approach taken in this study, including tailored intervention frequency and duration based on patient response, may serve as a model for future rehabilitation protocols aimed at similar patient populations.

**Table 1:** Results of symmetry for peak torque, work, and ROM.

CON/ECC			ECC/CON		
Symmetry of peak PF Torque	Pre-test	Post-test	Symmetry of peak PF Torque	Pre-test	Post-test
RI Peak TQ	0.30102	0.130102	RI Peak TQ	0.075	0.165517
RI Peak TQ/BW	0.300151	0.132331	RI Peak TQ/BW	0.079412	0.164969
RI Work	-1.36872	-2.70149	RI Work	-1.10127	0.202643
RI RoM	0.667347	0.358242	RI RoM	0.670213	0.655222
Symmetry of peak DF Torque	Pre-test	Post-test	Symmetry of peak DF Torque	Pre-test	Post-test
RI Peak TQ	0.270325	0.112621	RI Peak TQ	-0.36111	-0.46988
RI Peak TQ/BW	0.270983	0.113402	RI Peak TQ/BW	-0.36066	-0.46809
RI Work	-1.00881	-3.98492	RI Work	-2.44444	-0.6381

Note: RI; ratio index, TQ; torque, BW; body weight, RoM; range of motion.

**Table 2:** Results of symmetry for Area of center of pressure trajectory and path length.

Symmetry of Path Length	Pre-test	Post-test	Symmetry of Area	Pre-test	Post-test
Symmetry Index	-26779.9	-55948.8	Symmetry Index	-10600.2	-5680.35
Ratio Index	-0.04497	-0.2976	Ratio Index	-0.12459	-0.2976

**Table 3:** Results of symmetry for ankle passive stiffness in sagittal plane.

Symmetry of Stiffness R/L		Pre-test	Post-test
Symmetry Index		316800	12600
Ratio Index		-9	-7

**Table 4:** Results of symmetry for gait line length and contact time.

Symmetry of Gait Line Length	Pre-test	Post-test	Symmetry of Contact Time	Pre-test	Post-test
Symmetry Index	31338.4	17200.22	Symmetry Index	0.60175	0.28875
Ratio Index	0.14044	0.083441	Ratio Index	13.0631	4.104478

## Limitations and recommendation

Limitations of this case study include single case design reporting on a female patient with 30 years of regular exercise history, which may limit external validity. Inactive or overweight patients may have different outcomes. It's unclear whether IASTM or electrotherapy alone can lead to outcome improvements. Also, measuring asymmetries between sides is important to study injury-induced changes. Unilateral chronic ankle injury affects the kinematic parameters of the uninjured ankle and leads to morbidity on the contralateral side. Therefore, rehabilitation protocols should be planned on both sides. Non-weight-bearing dorsiflexion stiffness measures ankle flexibility differently than weight-bearing and may have different implications for injury potential. Future studies should measure stiffness during gait and jumping. Moreover, the participant could only perform the isokinetic test in 180 °/s. Thus, aerobic exercise should be added to the future plan.

## Conclusion

In conclusion, this case study demonstrated that a 12-week program consisting of IASTM and electrotherapy can have a positive impact on the symmetry of gait contact time, muscle strength, and ROM in a patient with Achilles tendon contracture. These findings have important implications for physical therapists and other healthcare providers, as they suggest that this combination of interventions may be an effective treatment approach for patients with similar conditions. However, further research is necessary to confirm these findings and to determine the optimal duration and intensity of treatment required to achieve these results.

## References

- Vieira Cardoso D, Dubois-Ferrière V, Gamulin A, Baréa C, Rodriguez P, Hannouche D, et al. Operatively treated ankle fractures in Switzerland, 2002-2012: epidemiology and associations between baseline characteristics and fracture types. *BMC Musculoskelet Disord.* 2021;22(1):266. Available from: <https://doi.org/10.1186/s12891-021-04144-5>
- Ramadi A, Beaupre LA, Heinrichs L, Pedersen ME. Recovery and Return to Activity 1 Year After Ankle Fracture Managed With Open Reduction and Internal Fixation: A Prospective Longitudinal Cohort Study. *Foot Ankle Orthop.* 2022;7(2):24730114221091806. Available from: <https://doi.org/10.1177/24730114221091806>
- Al Mahrouqi MM, MacDonald DA, Vicenzino B, Smith MD. Quality of life, function and disability in individuals with chronic ankle symptoms: a cross-sectional online survey. *J Foot Ankle Res.* 2020;13(1):67. Available from: <https://doi.org/10.1186/s13047-020-00432-w>
- Abbasi S, Mousavi SH, Khorramroo F. Association between lower limb alignment and low back pain: A systematic review with meta-analysis. *PLoS One.* 2024;19(10):e0311480. Available from: <https://doi.org/10.1371/journal.pone.0311480>
- Kaneguchi A, Ozawa J, Yamaoka K. The effects of immobilization duration on joint contracture formation after anterior cruciate ligament reconstruction in rats. *Clin Biomech (Bristol, Avon).* 2023;103:105926. Available from: <https://doi.org/10.1016/j.clinbiomech.2023.105926>
- Labidi M, Alhammoud M, Mtibaa K, Ihsan M, Deldicque L, Nasir N, et al. The Effects of Heat Therapy During Immobilization and Rehabilitation on Muscle Atrophy and Strength Loss at Return to Sports in Healthy Humans. *Orthop J Sports Med.* 2024;12(10):23259671241281727. Available from: <https://doi.org/10.1177/23259671241281727>
- Helme M, Tee J, Emmonds S, Low C. Does lower-limb asymmetry increase injury risk in sport? A systematic review. *Phys Ther Sport.* 2021;49:204-213. Available from: <https://doi.org/10.1016/j.ptsp.2021.03.001>
- Mousavi SH, Khorramroo F, Minoonejad H, Zwerver J. Effects of biofeedback on biomechanical factors associated with chronic ankle instability: a systematic review with meta-analysis. *BMC Sports Sci Med Rehabil.* 2023;15(1):168. Available from: <https://doi.org/10.1186/s13102-023-00780-7>
- Mousavi SH, Khorramroo F, Jafarnezhadgero A. Gait retraining targeting foot pronation: A systematic review and meta-analysis. *PLoS One.* 2024;19(3):e0298646. Available from: <https://doi.org/10.1371/journal.pone.0298646>
- Khorramroo F, Mousavi SH. The Effects of Minimal Shoes on Ankle Kinematics During Running: A Systematic Review. *J Adv Sport Technol.* 2022; 6:20-38. Available from: <https://doi.org/10.22098/jast.2022.1918>
- Khorramroo F, Mousavi SH, Minoonejad H. Lower Limb Coordination After Post-op Complications in a Case With Ankle Stiffness. *Phys Treat Phys Ther J.* 2024; 14:331-338. Available from: <https://ptj.uswr.ac.ir/article-1-632-en.html>
- Khorramroo F, Mousavi SH, Minoonejad H, Sorkhabi M. Managing Post-Operative Complications in a Patient with Excessive Ankle Stiffness: A Case Study Addressing Balance, Muscle Strength and Quality of Life. *Int J Mot Control Learn.* 2024; 5:e146340. Available from: <https://doi.org/10.5812/jmcl-146340>
- Torp DM, Thomas AC, Hubbard-Turner T, Donovan L. Biomechanical Response to External Biofeedback During Functional Tasks in Individuals With Chronic Ankle Instability. *J Athl Train.* 2021;56(3):263-271. Available from: <https://doi.org/10.4085/197-20>
- Kwon Y, Shin G. Foot kinematics and leg muscle activation patterns are altered in those with limited ankle dorsiflexion range of motion during incline walking. *Gait Posture.* 2022;92:315-320. Available from: <https://doi.org/10.1016/j.gaitpost.2021.12.002>
- Almutairi MK, Hunter GR, Lein DH, Kim S, Bryan DR, Inacio M, et al. Enhancement of Muscle Shortening Torque Preloaded with Muscle Lengthening is Joint-Specific. *J Hum Kinet.* 2023;87:11-21. Available from: <https://doi.org/10.5114/jhk/161729>
- Khorramroo F, Mousavi SH, Minoonejad H. Knee and Hip Kinematics during Gait following 12-week Conservative Interventions in a Case with Lower Limb Stiffness. *Int J Musculoskelet Pain Prev.* 2024; 9:1065-1069. Available from: <https://ijmpp.modares.ac.ir/article-32-75447-en.pdf>
- Kim YK, Cho SY, Lee KH. Effects of transcutaneous electrical nerve stimulation and instrument-assisted soft tissue mobilization combined treatment on chronic low back pain: A randomized controlled trial. *J Back Musculoskelet Rehabil.* 2021;34(5):895-902. Available from: <https://doi.org/10.3233/bmr-200369>
- Bush HM, Stanek JM, Wooldridge JD, Stephens SL, Barrack JS. Comparison of the Graston Technique® With Instrument-Assisted Soft Tissue Mobilization for Increasing Dorsiflexion Range of Motion. *J Sport Rehabil.* 2020;30(4):587-594. Available from: <https://doi.org/10.1123/jsr.2019-0397>
- Abdelatif EEM. Effect of transcutaneous electrical nerve stimulation and faradic current stimulation on the recovery of Bell's palsy. *Int J Hum Mov Sport Sci.* 2020; 8: 369-379. Available from: <https://www.hrpub.org/download/20201130/SAJ8-19920914.pdf>
- Chauhan IU, Telang PA. Effect of Myofascial Release (MFR) on Tendo Achilles (TA) Flexibility in Nurses: A Review. *Cureus.* 2022;14(11):e31319. Available from: <https://doi.org/10.7759/cureus.31319>



21. Kim DH, Lee JJ, Sung Hyun You J. Effects of instrument-assisted soft tissue mobilization technique on strength, knee joint passive stiffness, and pain threshold in hamstring shortness. *J Back Musculoskeletal Rehabil.* 2018;31(6):1169-1176. Available from: <https://doi.org/10.3233/bmr-170854>
22. Ganesh GS, Kumari R, Pattnaik M, Mohanty P, Mishra C, Kaur P, et al. Effectiveness of Faradic and Russian currents on plantar flexor muscle spasticity, ankle motor recovery, and functional gait in stroke patients. *Physiother Res Int.* 2018;23(2):e1705. Available from: <https://doi.org/10.1002/pri.1705>
23. Eagle SR, Kessels M, Johnson CD, Nijst B, Lovalekar M, Krajewski K, et al. Bilateral Strength Asymmetries and Unilateral Strength Imbalance: Predicting Ankle Injury When Considered With Higher Body Mass in US Special Forces<sup>a</sup>. *J Athl Train.* 2019;54(5):497-504. Available from: <https://doi.org/10.4085/1062-6050-255-18>
24. Gao Y, Arfat Y, Wang H, Goswami N. Muscle Atrophy Induced by Mechanical Unloading: Mechanisms and Potential Countermeasures. *Front Physiol.* 2018;9:235. Available from: <https://doi.org/10.3389/fphys.2018.00235>